


Known or Suspected TBI With Anticoagulant Use

MHealth Fairview System Guideline

Mechanism of Injury for potential TBI in a patient on Warfarin or Direct Oral Anticoagulants



*At Southdale Hospitalist will admit with Trauma or specialty consult

References:

1. Am Surg. 2017 May 1;83(5):429-435. Repeat Head CT? Not Necessary for Patients with a Negative Initial Head CT on Anticoagulation or Antiplatelet Therapy Suffering Low-Altitude Falls. Bauman ZM, Ruggero JM, Squindo S, McEachin C, Jaskot M, Ngo W, Barnes S, Lopez PP.
2. J Emerg Med. 2016 Nov;51(5):519-528. doi: 10.1016/j.jemermed.2016.05.045. Epub 2016 Jul 26. Risk of Delayed Intracranial Hemorrhage in Anticoagulated Patients with Mild Traumatic Brain Injury: Systematic Review and Meta-Analysis. Chauny JM, Marquis M, Bernard, Williamson D, Albert M, Laroche M, Daoust R.
3. J Neurol. 2018 Feb;265(2):315-321. doi: 10.1007/s00415-017-8701-y. Epub 2017 Dec 13. Evaluation of the yield of 24-h close observation in patients with mild traumatic brain injury on anticoagulation therapy: a retrospective multicenter study and meta-analysis. Verschoof MA, Zuurbier CCM, de Beer F, Coutinho JM, Eggink EA, van Geel BM.
4. https://www.facs.org/-/media/files/quality-programs/trauma/tqip/tbi_guidelines.ashx?la=en